

Astagon Tablet

Each Tablet contains:

Bromhexine8 mg
Chlorpheniramine Maleate.....4 mg
Ephedrine.....15 mg

COMPOSIT USES/ INDICATION OF ASTAGON :

Astagon Tablet - is indicated for **clinical relief of cough**

Bromhexine : Mucolytic agent used in the treatment of respiratory disorders associated with viscid or excessive mucus. Dry eye syndrome, Productive cough.

Chlorpheniramine Maleate : Acute urticaria, Advanced ovarian cancer, Allergic conjunctivitis, Atopic dermatitis, Chronic urticaria, Emergency treatment of anaphylactic reaction, Seasonal allergic rhinitis, Transcervical operative procedures in first trimester.

Ephedrine: Diabetic neuropathic edema, Narcolepsy, Nasal congestion, Nocturnal enuresis, Reduction of neutropenia-related risk of infection, Reversible airways obstruction.

DOSAGE AND ADMINISTRATION

Adults

Twice A Daily Or As Prescribe By Physician.

Children (6-12 years)

As Prescribe By Child Specialist.

COMPOSIT SIDE EFFECTS OF ASTAGON :

Bromhexine : Headache, GI disturbance, Skin Rash, Sweating, Dizziness, Bronchospasm, Rhinorrhoea, lacrimation, gastric irritation and hypersensitivity.

Chlorpheniramine Maleate : Jaundice, Sinus tachycardia, Seizures, Hypertension, Jaundice, CNS stimulation. Bone marrow suppression, Hypotension, Convulsions, Coma, Urinary retention, Unconsciousness, Fever, Dryness of mucous, Flushing membrane, Pupillary dilatation. Vomiting, Nausea, Sedation, Facial dyskinesias, Xerostomia, Abdominal pain, Constipation, Blurred vision, Dry mouth, Drowsiness, Restlessness, Palpitation, Dry mouth, Blood dyscrasias.

Ephedrine: Anxiety, tachycardia, tremor, dry mouth, hypertension, cardiac arrhythmias, impaired circulation to the extremities, nervousness, insomnia, palpitations. Difficulty in micturition in patients with prostatic enlargement. Cardiac arrhythmias.

PACKAGING INFORMATION

Astagon Tablets..... Blister Pack of 10 tablets