

Rovixim-Clav

www.greencrossindia.com

Cefixime and Clavulanate Potassium

COMPOSITION

ROVIXIM CLAV TABLETS

Each Film coated tablet contains:

Cefixime USP as trihydrate equivalent anhydrous cefixime200mg

Potassium Clavulanate Diluted IP equivalent to clavulanic acid125mg

Colour : Titanium dioxide IP

ROVIXIM CLAV DRY SYRUP

Each 5 ml suspension after reconstitution contains:

Cefixime IP as trihydrate equivalent anhydrous cefixime 50mg

Potassium Clavulanate Diluted IP equivalent to clavulanic acid ...31.25mg

INDICATIONS

ROVIXIM-CLAV is indicated for the treatment of :

Respiratory tract infections – Bronchitis, Bronchiectasis, Pneumonia

ENT Infections – Chronic Maxillary Sinusitis, Chronic Otitis Media

Urinary tract infections – Acute uncomplicated and complicated urinary tract infection

DOSAGE AND ADMINISTRATION

Adults and Children over 10 Years : One tablet twice daily

The usual course of treatment is 7 days. This may be continued for up to 14 days if required.

CONTRAINDICATIONS

Patients with known hypersensitivity to cephalosporin antibiotics.

PACKAGING INFORMATION

ROVIXIM-CLAV – Each Strip contains 6 tablets